

# Chapter 9: Independence and Conditioning (Joint Distributions)

Meike Niederhausen and Nicky Wakim

2023-10-11

# Table of contents

- Learning Objectives
- What is a joint pmf?
- This chapter's main example
- Joint pmf
- Marginal pmf's
- Remarks on the joint pmf
- What is a joint CDF?
- Joint CDFs
- Marginal CDFs
- Remarks on the joint and marginal CDF
- Independence and Conditioning
- What is the conditional pmf?
- Remarks on the conditional pmf
- Conditional pmf's

# Learning Objectives

- 1. Calculate probabilities for a pair of discrete random variables
- 2. Calculate and graph a *joint, marginal, and conditional* probability mass function (pmf)
- 3. Calculate and graph a *joint, marginal, and conditional* cumulative distribution function (CDF)

# What is a joint pmf?

## Definition: joint pmf

The **joint pmf** of a pair of discrete r.v.'s  $X$  and  $Y$  is

$$p_{X,Y}(x, y) = \mathbb{P}(X = x \text{ and } Y = y) = \mathbb{P}(X = x, Y = y)$$

$$p_X(x) \quad p_Y(y)$$

# This chapter's main example

## Example 1

Let  $X$  and  $Y$  be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

- 1. Find  $p_{X,Y}(x, y)$ .
- 2. Find  $\mathbb{P}(X + Y = 3)$ .
- 3. Find  $\mathbb{P}(Y = 1)$ .
- 4. Find  $\mathbb{P}(Y \leq 2)$ .
- 5. Find the joint CDF  $F_{X,Y}(x, y)$  for the joint pmf  $p_{X,Y}(x, y)$
- 6. Find the marginal CDFs  $F_X(x)$  and  $F_Y(y)$
- 7. Find  $p_{X|Y}(x|y)$ .
- 8. Are  $X$  and  $Y$  independent? Why or why not?

# Joint pmf

## Example 1

Let  $X$  and  $Y$  be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

1. Find  $p_{X,Y}(x,y)$ .

2. Find  $\mathbb{P}(X + Y = 3)$ .

$X$   
(1st  
draw)

	$Y$ (2nd draw)		
	1	2	3
1	0	$\frac{1}{6}$	$\frac{1}{6}$
2	$\frac{1}{6}$	0	$\frac{1}{6}$
3	$\frac{1}{6}$	$\frac{1}{6}$	0

drawing 1 ( $x$ ) then 1 ( $y$ )  
is impossible b/c no replacement

$$P_{X,Y}(x=1, y=1) = 0$$

$$P_{X,Y}(x=1, y=2) = \left(\frac{1}{3}\right)\left(\frac{1}{2}\right) = \frac{1}{6}$$

$$P_{X,Y}(x=1, y=3) = \left(\frac{1}{3}\right)\left(\frac{1}{2}\right) = \frac{1}{6}$$

# Marginal pmf's

## Example 1

Let  $X$  and  $Y$  be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

3. Find  $\mathbb{P}(Y = 1)$ .

4. Find  $\mathbb{P}(Y \leq 2)$ .

# Remarks on the joint pmf

Some properties of joint pmf's:

- A joint pmf  $p_{X,Y}(x, y)$  must satisfy the following properties:
  - $p_{X,Y}(x, y) \geq 0$  for all  $x, y$ .
  - $\sum_{\{\text{all } x\}} \sum_{\{\text{all } y\}} p_{X,Y}(x, y) = 1$ .
- Marginal pmf's:
  - $p_X(x) = \sum_{\{\text{all } y\}} p_{X,Y}(x, y)$
  - $p_Y(y) = \sum_{\{\text{all } x\}} p_{X,Y}(x, y)$



# What is a joint CDF?

## Definition: joint CDF

The **joint CDF** of a pair of discrete r.v.'s  $X$  and  $Y$  is

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x \text{ and } Y \leq y) = \mathbb{P}(X \leq x, Y \leq y)$$

# Joint CDFs

## Example 1

Let  $X$  and  $Y$  be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

5. Find the joint CDF  $F_{X,Y}(x, y)$  for the joint pmf  $p_{X,Y}(x, y)$

# Marginal CDFs

## Example 1

Let  $X$  and  $Y$  be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

6. Find the marginal CDFs  $F_X(x)$  and  $F_Y(y)$

# Remarks on the joint and marginal CDF

- $F_X(x)$ : right most columns of the Cdf table (where the  $Y$  values are largest)
- $F_Y(y)$ : bottom row of the table (where  $X$  values are largest)
- $F_X(x) = \lim_{y \rightarrow \infty} F_{X,Y}(x, y)$
- $F_Y(y) = \lim_{x \rightarrow \infty} F_{X,Y}(x, y)$

# Independence and Conditioning

Recall that for events  $A$  and  $B$ ,

- $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- $A$  and  $B$  are independent if and only if
  - $\mathbb{P}(A|B) = \mathbb{P}(A)$
  - $\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$

Independence and conditioning are defined similarly for r.v.'s, since

$$p_X(x) = \mathbb{P}(X = x) \text{ and } p_{X,Y}(x, y) = \mathbb{P}(X = x, Y = y).$$

# What is the conditional pmf?

## Definition: conditional pmf

The **conditional pmf** of a pair of discrete r.v.'s  $X$  and  $Y$  is defined as

$$p_{X|Y}(x|y) = \mathbb{P}(X = x|Y = y) = \frac{\mathbb{P}(X = x \text{ and } Y = y)}{\mathbb{P}(Y = y)} = \frac{p_{X,Y}(x, y)}{p_Y(y)}$$

if  $p_Y(y) > 0$ .

# Remarks on the conditional pmf

The following properties follow from the conditional pmf definition:

- If  $X \perp Y$  (independent)
  - $p_{X|Y}(x|y) = p_X(x)$  for all  $x$  and  $y$
  - $p_{X,Y}(x, y) = p_X(x)p_Y(y)$  for all  $x$  and  $y$
  - Which also implies ( $\Rightarrow$ ):  $F_{X,Y}(x, y) = F_X(x)F_Y(y)$  for all  $x$  and  $y$
- If  $X_1, X_2, \dots, X_n$  are independent

- $$p_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) = \prod_{i=1}^n p_{X_i}(x_i)$$

- $$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n) = \prod_{i=1}^n P(X_i \leq x_i) = \prod_{i=1}^n F_{X_i}(x_i)$$

# Conditional pmf's

## Example 1

Let  $X$  and  $Y$  be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

7. Find  $p_{X|Y}(x|y)$ .

8. Are  $X$  and  $Y$  independent? Why or why not?

Remark:

- To show that  $X$  and  $Y$  are *not* independent, we just need to find one counter example
- However, to show that they *are* independent, we need to verify this for all possible pairs of  $x$  and  $y$



